

**Nebraska Soybean Board**  
**Year-End Research Findings Report**



10/31/2017

Please use this form to summarize the practical benefits of your research project and what has been accomplished. Your answers need to convey why the project is important and how the results impact soybean production.

**Project Title:** *Benchmarking soybean production systems in Nebraska (#90005)*

**Contractor & Principal Investigator:** *UNL—Department of Agronomy. PI: Patricio Grassini*

Please check/fill in appropriate box:  Continuation research project  
 Year 2 of 2 research project (for example: Year 1 of 2)

**1. What was the focus of the research project or educational activity?**

*The primary goal of the proposed project is to “benchmark” current yield and management practices in producer fields. This project is a sub-component of a larger, regional 10-state benchmarking project led by PI Grassini and supported by the North Central Soybean Research Program (NCSRP). NSB funding contributed to data collection in Nebraska.*

**2. What are the major findings of the research or impacts of the educational activity?**

*We partnered with 20 UNL Extension Educators and NE Natural Resources Districts (NRDs) to collect the data. The number of filled surveys collected by UNL Extension Educators, together with the surveys filled out by NRD soybean producers, sum up to 414 soybean fields in 2014, 483 soybean fields in 2015, and 642 soybean yields in 2016 (total of 1,539 fields over the 2-year project). Note that the number of surveyed soybean fields is almost four times the target number (480 fields) set at the beginning of the project. Relative to other states that participate in the NCSRP-funded project, the largest number of survey forms was collected from Nebraska thanks to the help of Nebraska Extension and NRDs and the support from the NSB. So, we are very happy on how well the collaboration with UNL extension and NRD worked out and we look forward to collecting the data following the same model during the next winter! The core team at UNL have inputted, quality control, and archived the data collected by Extension Educators, NRDs, etc. in a digital database. Weather and soil data were retrieved for each individual field, which will allow proper contextualization of the collected data. A detailed report summarizing the collected data has been prepared (see appended TECHNICAL REPORT). Interesting findings based on collected data are:*

- *Nebraska average dryland and irrigated yields were 56 bu/ac and 67 bu/ac respectively, both above the average soybean yield in the north-central region (54 bu/ac). Only a small proportion of producers (2%) attained soybean yields near or above 80 bu/ac.*
- *Half of the soybean area in the north-central region is no-till. Adoption of no-till in Nebraska is greater in dryland (77% of fields) than in irrigated fields (51% of fields).*
- *About 25% of soybean fields in this region are planted during the first week of May or earlier. This figure rises to 45% in Nebraska.*
- *Seeding rates used by producers are well above economically optimal soybean seeding rates, which, in the case of Nebraska, is 120,000 seeds/acre.*
- *Most producers in the region grow soybean at a 15-inch row spacing, except for Nebraska and eastern Iowa where 30-inch spacing still prevails.*
- *Across the entire north-central region, 8%, 19% and 24% of soybean fields are treated with foliar fungicide only, insecticide only, and both fungicide and insecticide, respectively. In Nebraska, these figures are lower at 6%, 3% and 17%.*

**\*\*This form must be completed and submitted with the fourth quarter report.**

## **Nebraska Soybean Board Year-End Research Findings Report**

*Please use this form to summarize the practical benefits of your research project and what has been accomplished. Your answers need to convey why the project is important and how the results impact soybean production.*

**- About 15%, 54% and 54% of fields in the north-central region received starter, phosphorous, and potassium fertilizer, respectively. In Nebraska, these figures are 15%, 70%, and 54%**

**3. Briefly summarize, in lay terms, the impact your findings have had, or will have, on improving the productivity of soybeans in Nebraska and the U.S.**

***We have compiled the most extensive, detailed, and agronomically-relevant database on soybean production systems in USA and worldwide. For first time, it is possible to examine spatial variation (across and within states) in soybean yield and management practices. This information will be very useful at determining the factors that can help increase soybean yield, input-use efficiency, or both in producer fields in Nebraska and the rest of the US north-central region and, in doing so, increase on-farm net profit.***

**4. Describe how your findings have been (or soon will be) distributed to (a) farmers and (b) public researchers. List specific publications, websites, press releases. etc.**

***A summary report was prepared and posted in the North Central Soybean Research Program website (<http://www.soybeanresearchinfo.com/index.php?id=57> -- **SEE APPENDED TECHNICAL REPORT**) and also published as a UNL CropWatch article (<http://cropwatch.unl.edu/2016/help-us-identify-limiting-factors-nebraska-soybean-fields>) and made accessible to all NE soybean producers. The report was also shared with educators, NRDs, NSB members, etc. Likewise, two scientific paper were published derived from the data collected for this project:***

***Mourtzinis S, Rattalino Edreira JI, Conley SP, Grassini P (2017) From grid to field: assessing quality of gridded weather data for agricultural applications. European J. of Agronomy 82, 163-172.***

***Edreira JI, Mourtzinis S, Conley SP, Roth A, Ciampitti IA, Licht MA, Kandel H, Kyveryga PM, Lindsey LE, Mueller DS, Naeve SL, Nafziger E, Specht JE, Stanley J, Staton MJ, Grassini P (2017) Assessing causes of yield gaps in agricultural areas with diversity in climate and soils. Agricu. For. Meteorol. 247, 170-180.***

***Likewise, Patricio Grassini (Project PI) gave presentations at the 2016 & 2017 winter UNL Crop Production Clinics at 8 locations in NE, and will also be presenting results during the 2018 clinics. Also, Juan Ignacio Rattalino Edreira (Post-Doctoral Research Associate working on this project) presented results from this project at the 2016 Annual ASA/CSSA/SSSA Meetings at Phoenix AZ and will present again at the 2017 meetings at Tampa FL.***

**5. Did the NE soybean checkoff funding support for your project leverage any additional state or Federal funding support? (Please list sources and dollars approved.)**

***The NSB funding helped us to secure funding from North Central Soybean Research Program (NCSRP) to continue the regional (10-state) benchmarking project for another year (year 3), at a total level of 1.5 million (total for the 3 years). Patricio Grassini is the PI of the funded NCSRP project.***

**SEE APPENDED TECHNICAL REPORT.**

***\*\*This form must be completed and submitted with the fourth quarter report.***

# Key Management Practices That Explain Soybean Yield Gaps Across the North Central US

**Juan I. Rattalino Edreira,  
James E. Specht, Patricio Grassini**  
Department of Agronomy and Horticulture,  
University of Nebraska-Lincoln

**Spyridon Mourtzinis, Shawn P. Conley,  
Adam C. Roth**  
Department of Agronomy,  
University of Wisconsin-Madison

**Ignacio A. Ciampitti**  
Department of Agronomy,  
Kansas State University

**Mark A. Licht**  
Department of Agronomy,  
Iowa State University

**Hans Kandel, Jordan Stanley**  
Department of Plant Sciences,  
North Dakota State University

**Peter M. Kyveryga**  
Iowa Soybean Association

**Laura E. Lindsey**  
Department of Horticulture and Crop Science,  
Ohio State University

**Daren S. Mueller**  
Department of Plant Pathology and Microbiology,  
Iowa State University

**Seth L. Naeve**  
Department of Agronomy and Plant Genetics,  
University of Minnesota

**Emerson Nafziger**  
Department of Crop Sciences, University of Illinois

**Michael J. Staton**  
Michigan State University Extension

## Highlights


- ▶ We developed a novel approach that combines producer survey data with a biophysical spatial framework for identifying causes of yield gaps over large agricultural areas with diversity in climate and soils.
- ▶ The approach was applied to both rainfed and irrigated soybean in the North Central US region, and it was based on producer survey data on yield and management collected from 3,568 fields over two crop seasons.
- ▶ The analysis indicated that the average regional yield potential was 71 bu ac<sup>-1</sup> (rainfed) and 85 bu ac<sup>-1</sup> (irrigated), with a respective yield gap of 22% and 13% of maximum yield potential.
- ▶ Planting date, tillage, and in-season foliar fungicide and/or insecticide were identified as explanatory causes for yield variation, with planting date the most consistent management factor that influenced soybean yield.

## Introduction

To date identification of causes of yield gaps (difference between maximum yield potential and measured yield in producer yields) has been restricted to small geographic areas. In this study, we developed a novel approach that combines producer-reported data and a spatial framework to identify explanatory causes of yield gap over large geographic regions with diversity of climate, soils, and water regimes (rainfed and irrigated). We focused on soybean in the North-Central United States region, which accounts for approximately one third of global soybean production, as a case study to provide a proof of concept on the proposed approach. The specific objectives of this project were to evaluate the proposed approach for its ability to: (1) benchmark producer soybean yields in relation to yield potential of their fields, (2) identify key management practices that explain yield gaps, and (3) explain the drivers for some of the observed (M)anagement × (E)nvironment interactions.

## Producer data collection and quality control

Data on soybean yield and management practices were collected over two crop seasons (2014 and 2015) from fields planted to soybean in 10 states in the North Central US region: Illinois (IL), Indiana (IN), Iowa (IA), Kansas (KS),



**Figure 1.** Example of an actual survey form from a Nebraska soybean producer that provides information for three irrigated fields and one rainfed field planted to soybean in 2014 and 2015. This survey form was used to collect information from producer fields across 10 states in the North Central US region. Note that producer name is not shown and field location was hatched in order to keep personal information confidential.

**PRODUCER NAME:** [REDACTED] **MAILING ADDRESS:** [REDACTED]

Please provide information for four SOYBEAN fields on your farm in 2014. If you have questions, contact Professor **Patricio Grassini** (Phone: 402-472-5554 / e-mail: pgrassini2@unl.edu). Note that all provided info will be kept confidential! An **EXAMPLE** is shown in red.

	EXAMPLE	2014 Soybean	2014 Soybean	2015 Soybean	2015 Soybean
Specify field location by Section: Township: Range: →	NE 1/4 29N 28W				
Please sketch-in the boundaries of your field location within the Section					
OR GPS coordinates of field centroid:	41.678, -100.257	[REDACTED]	[REDACTED]	[REDACTED]	[REDACTED]
OR County & field location relative to Rd Intersection:	Saunders Co. SW of Rd 11 & N	[REDACTED]	[REDACTED]	[REDACTED]	[REDACTED]
Dryland? OR Pivot, Gravity? indicate field size (acres)	Pivot (130 ac)	Pivot (137 ac)	Grassy (30 ac)	Pivot (84 ac)	Dryland (11 ac)
Does this field have drainage? (no, old clay tile, new systematic tile, surface drainage, other)	No	No	No	No	No
Total Inches of Irrigation Applied to crop?	5 inches	3.5 in.	4.5 in.	3.5 in.	0
SOYBEAN YIELD (bushels/acre) for this FIELD:	70	80	70	85	89
Lowest   Highest Yield (bu/ac) of your soy fields that year	Low: 62 High: 80	Low: 61 High: 90	Low: 55 High: 86	Low: 61 High: 90	Low: 13 High: 66
*Use Irrigated fields yield range if this crop was Irrigated: *Use Dryland fields yield range if this crop was Dryland:					
Planting Date in this FIELD (Month/Day/Year):	5/15/2014	4/25/2014	5/2/2014	4/25/2014	5/14/2014
Variety Name (Brand & Number):	Pioneer P93M11	Chancellor RR2	Chancellor RR2	Chancellor RR2	Chancellor
Seeding Rate (seeds/ac):	125,000	140,000	140,000	140,000	140,000
Row spacing (inches):	30	30	30	15	15
Seed Treated (Yes/No)? What Brand Name Product(s)?	Yes (Cruiser-Max)	yes Ardelexon	yes Ardelexon	yes Ardelexon	yes Ardelexon
Prior Crop in this FIELD? Residue harvested or grazed?	Corn - Grazed	Corn - Grazed	Corn - No	Corn - Grazed	Corn - No
Tillage after prior crop? No-Till (NT); Ridge (RT); Strip (ST); Disk (D); Chisel (C); Vertical (V) - indicate timing (month-year)	ST (March-2014)	NT	D (April 2014)	NT	NT
Any (non-starter) fertilizer after prior crop?	P:0; K:0; S:30	P:0; K:0	P:0; K:0	P:0; K:0	P:0; K:0
Specify rate (pounds NUTRIENT/ac) and timing (month-year)	Other: S (11 lbs) Time: March-2014	Other: None Time: None			
Any STARTER fertilizer (Yes/No)? If Yes, specify nutrients	Yes (N, P, Zn)	No	No	No	No
Any Lime (L) or Manure (M)? If yes, specify timing (mm-yy)	M (Nov-2013)	No	No	No	No
PRE- or POST-emergence herbicide program or BOTH?	Both	Both	Both	Both	Both
Any in-season foliar fungicide (F) / insecticide (I)?	F and I	No	No	No	No
Soy Cyst Nematodes (Yes/No/I don't know)?	No	No	No	No	No
Iron Deficiency Chlorosis (Yes/No)?	No	No	No	No	No
Any significant yield loss due to insects, Diseases, Weeds, Frost, Hail, Flood, Lodging? Specify problem	Frost (Sept-2014)	None	None	None	Hail (July 2014)

Michigan (MI), Minnesota (MN), Ohio (OH), Nebraska (NE), North Dakota (ND), and Wisconsin (WI). Soybean producers provided data via returned surveys distributed by local crop consultants, Extension educators, soybean grower boards, and Natural Resources Districts (Figure 1). Briefly, producers were asked to report the range of average field yield across the fields planted to soybean in each year and water regime and to provide data for a number of fields that portray well that yield range. Requested data also included field location, average field yield, crop management (e.g., planting date, seeding rate, row spacing, cultivar, and tillage method), applied inputs (e.g., irrigation, fertilizer, lime, manure, and pesticides), and incidence of biotic and abiotic factors (e.g., insect pests, diseases, weeds, hail, waterlogging, and frost). Survey data were inputted into a digital database and screened to remove erroneous or missing data entries. We were interested in yield variation as related with management factors; hence, a few fields with extremely low yield due to incidence of unmanageable production site adversities (hail, waterlogging, wind, and frost) were excluded from the analyses. After quality control, the database contained data from a total of 3,216 fields planted to soybean in 2014 and 2015.


**Producer data stratification based on soil-climate conditions**

A major challenge with this kind of data is how to cluster producer fields in order to identify management factors that consistently lead to higher yields for a given climate-soil combination. In the present study, surveyed fields were grouped based upon their climate and soil using the spatial framework developed for the central and eastern US by the Global Yield Gap Atlas (<http://www.yieldgap.org>). This framework delineates regions [hereafter called technology extrapolation domains (TEDs)] based on four biophysical attributes

**Figure 2.** Map of the North Central US region showing nine technology extrapolation domains (TEDs) and meteorological stations (solid circles) selected for the present study. A coding system (from TED 1 to 9) is used to identify each TED (shown with a unique color) and its associated water regime (I: irrigated, R: rain-fed). There were actually 10 TED-water regimes (denominated as just TEDs for simplicity) because rainfed and irrigated fields co-existed in TED 7 (7R and 7I, respectively).

**Top inset.** Soybean harvested area in year 2015 (green area; USDA-NASS, 2016b) and location of the 3,216 surveyed soybean fields (red dots).


**Bottom inset.** Location of North Central US region — 12 states within the conterminous US.



that govern crop yield and its inter-annual variability: (1) annual total growing degree-days, which, in large part, determines the length of crop growing season, (2) aridity index, which largely defines the degree of water limitation in rainfed cropping systems, (3) annual temperature seasonality, which differentiates between temperate and tropical climates, and (4) plant-available water holding capacity in the rootable soil depth, which determines the ability of the soil to supply water to support crop growth during rain-free periods. We selected TEDs that portrayed the diversity of climate, soils, and water regimes in the North Central US region (Figure 2). Six TEDs included only rainfed soybean fields (1R, 2R, 3R, 4R, 5R, and 6R) while two TEDs included only irrigated soybean fields (8I and 9I). One TED included both irrigated and rainfed soybean fields (7I and 7R). Because the impact of management factors on yield is influenced by water supply, we separated water regimes (WR; rainfed and irrigated) within the same TED. Hence, a total of 10 TED-WR combinations were eventually used in this study, which are referred hereafter as TEDs for simplicity (total of 10 TEDs). Selected TEDs included 38% of the surveyed fields (total of 1343 fields) and accounted for 25 and 45% of US rainfed and irrigated soybean area, respectively. Each individual TED contained  $\geq 98$  (rainfed) and  $\geq 59$  (irrigated) surveyed fields, with an average of 137 fields per TED.

### Yield potential, average producer yield, and yield gaps

Annual yield potential ( $Y_p$ , yield potential of irrigated field) and water-limited yield potential ( $Y_w$ , yield potential of rainfed fields) were estimated using measured daily weather data (including solar radiation, rainfall, and maximum and minimum air temperature) collected at 2–3 meteorological stations located within each TED, preferably in proximity to the areas with the highest density of surveyed fields.  $Y_w$  and  $Y_p$  were used as benchmarks for calculating yield




gap for rainfed (TEDs 1R, 2R, 3R, 4R, 5R, 6R, and 7R) and for irrigated TEDs (7I, 8I, and 9I). The yield gap was calculated as the difference between  $Y_p$  (or  $Y_w$ ) and average producer yield and expressed as percentage of  $Y_p$  (irrigated) or  $Y_w$  (rainfed).

Average  $Y_w$  ranged from 48–80  $\text{bu ac}^{-1}$ , while  $Y_p$  varied from 80–91  $\text{bu ac}^{-1}$  across TEDs (Figure 3). TED 3R exhibited the lowest  $Y_w$  due to lower seasonal precipitation in relation with other TEDs. In contrast,  $Y_p$  was highest in TED 8I due to non-limiting water supply and high incident solar radiation. Upscaled to the entire North Central US region,  $Y_w$  and  $Y_p$  averaged 71 and 85  $\text{bu ac}^{-1}$ , respectively. Average producer yield was consistently lower than  $Y_w$  (or  $Y_p$ ) across all TEDs ( $p < 0.01$ ), and there was a large variation in average annual yield across TEDs, ranging from 39–73  $\text{bu ac}^{-1}$ . Yield gap, expressed as percentage of  $Y_p$  (irrigated) or  $Y_w$  (rainfed), tended to be larger in rainfed (range: 15–28%) than in irrigated TEDs (range: 11–16%). At the regional level, the rainfed yield gap averaged 22% in contrast to the irrigated yield gap of 13%.

### Management practices explaining yield gap between high- and

**Figure 3.** Yield potential for rainfed ( $Y_w$ ) and irrigated ( $Y_p$ ) soybean in each of the 10 TEDs in 2014 (14) and 2015 (15). Solid and empty portions of the bars represent the average producer yield and yield gap, respectively. Values on top of the bars indicate the (2-year) average yield gap, expressed as percentage of  $Y_w$  (rainfed) or  $Y_p$  (irrigated).




### low-yield fields

As a first approach to identify factors explaining yield gap, high-yield (HY) and low-yield (LY) field classes were identified based on their respective presence in the upper and lower terciles (top 1/3 versus bottom 1/3 of fields) of the field yield distribution within each TED. Analysis of management practices allowed identification of candidate factors explaining yield gap in each TED. Differences in planting date, tillage, in-season foliar fungicide and/or insecticide, drainage system, and soybean cultivar maturity group (MG) between high- and low-yield fields were statistically significant in half or more of the 10 TEDs ( $p < 0.10$ ).

### Planting date: The main explanatory factor

Planting date had the most consistent impact on soybean yield (Figure 4), representing 28% of the total yield gap across TEDs (range: 2–56%). HY fields were sown, on average, 7 days earlier than LY fields in both irrigated and

**Figure 4.** Producer soybean yield plotted against planting date in 10 technology extrapolation domains (TED) in the NC USA region, including rainfed (A–G) and irrigated (G–I) production areas. Solid line corresponds to the fitted boundary function using quantile regression (percentile 90th). Separate boundaries were derived for rainfed (empty symbols) and irrigated (solid symbols) soybean fields in TED7. Slope of the fitted boundary function ( $b$ ) is shown, with asterisks indicating significance at  $p < 0.1^*$ ,  $p < 0.05^{**}$ , and  $p < 0.01^{***}$  for the null hypothesis of  $b = 0$ .




rainfed conditions. There was a strong planting date  $\times$  TED interaction on yield as indicated by the wide range in yield penalty across TEDs, ranging from 0 to  $-0.5 \text{ bu ac}^{-1} \text{ day}^{-1}$  (Figure 4).

Assessment of the observed TED  $\times$  M interactions, in relation to weather dynamics during the growing season, revealed a relationship between yield response to planting date and the degree of water deficit during pod setting (R3–R5) phase (Figure 5). Yield penalty (or response) to planting date was negligible when water balance was  $< -4$  inches, but increased linearly up to nearly  $-1.6$  inches. Yield response to planting date remained relatively unchanged at water balance  $> -1.6$  inches, ranging from  $0.3$ – $0.5 \text{ bu ac}^{-1} \text{ day}^{-1}$ . The role of water balance in influencing the yield response to planting date was evident for TED 7, where irrigated and rainfed crops exhibited a six-fold difference ( $0.5$  versus  $0.1 \text{ bu ac}^{-1} \text{ day}^{-1}$ , respectively) (Figure 4). In other words, these findings indicated that yield response to planting date diminished as the degree of water limitation in the pod-setting period of the production environment increases. It was notable that yield response to planting date delay exhibited much higher explanatory power with the degree of water deficit during pod setting phase

01) relative to the other crop phases (early vegetative phase, late vegetative phase, and seed filling) or entire crop season ( $r^2 < 0.38$ ,  $p > 0.06$ ).

**Figure 5.** Soybean yield penalty due to planting date delay as a function of water balance during the pod-setting (R3–R5) phase across 10 technology extrapolation domains (TEDs) including rainfed (yellow circles) and irrigated (blue circles) production environments (averaged over 2014–2015). Water balance was estimated as the difference between rainfall and simulated non-water limiting crop evapotranspiration and set at zero for irrigated crops. Parameters of the fitted linear-plateau model (solid line) and coefficient of determination ( $r^2$ ) are shown.




### Tillage, fungicide and/or insecticide applications, drainage system, and soybean maturity groups

Similarly to planting date, other management practices also exhibited a significant M × TED interaction (Figure 6). For this analysis, fields were categorized as either no-till or tilled, with the latter including chisel, disk, strip-till, ridge-till, vertical, field cultivator, and moldboard plow. We did not find evidence of no-till fields outperforming yield of tilled fields in every TED; indeed, tilled fields yielded significantly more in half of the TEDs (2.3 bu ac<sup>-1</sup>;  $p = 0.02$ ) (Figure 6). However, there may still be other functional reasons for producers to adopt no-till despite the observed yield penalty. For example, no-till can help control soil erosion and reduce irrigation water requirements. Indeed we found that, on average, total irrigation was 2.5 inches less in no-till versus tilled fields ( $p < 0.01$ ).

While there was an overall statistically positive impact of foliar fungicide and/or insecticide (4.6 bu ac<sup>-1</sup>,  $p < 0.01$ ) and artificial drainage (2.7 bu ac<sup>-1</sup>;  $p = 0.05$ ) on soybean seed yield, the magnitude of these yield differences were not consistent across TEDs and not even significant in some of them (Figure 6). For example, average yield of fields treated with foliar fungicide and/or insecticide was 11.2 bu ac<sup>-1</sup> higher in relation with untreated fields in TED 7R, but this yield difference was negligible (-0.9 bu ac<sup>-1</sup>) and not statistically significant in TED 6R. Likewise, artificially drained fields achieved statistically higher yields compared with fields without artificial drainage in only two of six TEDs. Although differences in variety MG between high- and low-yield fields were less than one unit, there was a consistent trend towards shorter MGs in the high-yield field tercile (top 1/3) in all TEDs, except for those located in the northern fringe of the North Central US region (3R and 4R).



**Figure 6.** Comparison of average producer soybean yield between groups of fields with different management practices across ten technology extrapolation domains (TEDs): (A) tillage (tilled versus no-till), (B) in-season foliar fungicide and/or insecticide (treated versus untreated fields), and (C) artificial drainage (fields with and without artificial drainage system). Star inside symbols indicate statistically significant difference for a given TED (t-test;  $p < 0.1$ ). Asterisks indicate significance of the impact on yield with respect to the specified management factor (M), and its interaction with year (M  $\times$  Y) or with TED (M  $\times$  TED) as evaluated using F-test at  $p < 0.1$  (\*),  $p < 0.05$  (\*\*), and  $p < 0.01$  (\*\*\*)). Data from the two crop seasons were pooled for the analysis because M  $\times$  Y influence on yield was not statistically significant. TEDs 7R, 7I, 8I, and 9I are not included in (C) because of the low number of fields with artificial drainage.




### Other management factors with low influence on yield gap

In contrast to the aforementioned variables, there were inconsistent (and generally small) differences between HY and LY fields in relation to row spacing, seeding rate, seed treatment, nutrient (N, P, K) fertilizer application, lime, and manure. Lack of statistically significant differences between management practices need to be interpreted with caution. For example, some practices might influence yield depending upon the level of another management practice [e.g., seed treatment in relation with planting date (Gaspar and Conley, 2015)]. Likewise, the benefit of other practices may only be realized in crop seasons with unfavorable weather, which was not the case in our study [e.g., narrow row spacing, no-till (Taylor, 1980; Wilhelm and Wortmann, 2004)]. Similarly, yield impact of some practices may be masked by other field variables not accounted here. For example, lack of yield differences between fields that received fertilizer application versus those that did not receive fertilizer might reflect producer tendency to apply fertilizer only in fields where soil nutrient status is inadequate as evaluated using soil nutrient tests. It may also reflect that many producers over-fertilized the previous corn crop expecting the subsequent soybean crop to benefit from the residual soil fertility. Finally, there are management practices that exhibited a very narrow range (e.g., MG) or inputs that are applied in amounts well above their optimums. For example, on-farm average soybean seeding rate ranged from 147,000 to 172,000 seeds  $\text{ac}^{-1}$  across TEDs. These densities are higher than the required plant density for maximum yields (100,000–145,000 plants  $\text{ac}^{-1}$ ) (Grassini et al., 2015); hence, our analysis will not fully capture the influence of these management factors on soybean yield.

### Final consideration

Beside the identification of yield gap causes, another contribution of the present study is to provide a solid basis to assess what would be the extra crop production, at both local (TED) and regional (North Central US) levels, that would



### Acknowledgements

The authors would like to thank the North-Central Soybean Research Program, Nebraska Soybean Board, and Wisconsin Soybean Marketing Board for their support to this project, and the University of Nebraska-Lincoln Extension Educators, Nebraska Natural Resource Districts, and Iowa Soybean Association for helping collect the producer data.

Finally, we would like to thank Lim Davy, Agustina Diale, Laurie Gerber, Clare Gietzel, Mariano Hernandez, Ngu Kah Hui, Caleb Novak, Juliana de Oliveira Hello, Matt Richmond, and Paige Wacker for inputting and cleaning the survey data.

result from complete producer adoption or fine-tuning of a given management practice. For example, the potential extra production derived from earlier soybean planting can be calculated based on the (1) specific yield response to planting date in each TED, (2) the degree to which the current average planting date differs from the optimal one, and (3) soybean harvested area in each TED. For example, a 2-week shift towards early soybean planting in TED 4R, from current average planting on May 17 to a hypothetical, yet realistic, May 3 planting, would result in 5.2 bu ac<sup>-1</sup> yield increase and 18.5 million bu production increase, leading to a 10% and 0.7% increase in soybean production in TED 4 and North Central US region, respectively. This example illustrates the power of this approach for impact assessment to support policy and investment prioritization and for monitoring the impact of research and Extension programs.

### Conclusion

Soybean yield gap and its causes were assessed for the North Central US region using a novel approach that combines a spatial framework and producer self-reported data. The framework applied in this study explained the largest portion of the spatial variation in yield and management practices across the North Central US region. Soybean yield gap in the North Central US were relatively small, averaging 22% (rainfed) and 13% (irrigated) of the estimated yield potential. Planting date was the most consistent factor explaining yield variation within the same TED and year, with magnitude of yield response to planting delay dependent upon degree of water deficit during pod setting phase. Other practices also explained yield variation (tillage, and in-season foliar fungicide and/or insecticide, and artificial drainage), but the degree to which each of these practices influences yield depended upon TED. The combined use of producer data and a robust spatial framework that captured regional variation in weather and soils represents a cost-effective approach to identify causes of yield gap across large geographic regions, which, in turn, can help inform and strategize research and Extension programs at both local and regional levels.

### References

- Gaspar A.P., Conley S.P. (2015) *Responses of canopy reflectance, light interception, and soybean seed yield to replanting suboptimal stands*. *Crop Sci* 55:377-385. DOI: 10.2135/cropsci2014.03.0200.
- Grassini P., Specht J.E., Tollenaar M., Ciampitti I.A., Cassman K.G. (2015) *High-yield maize-soybean cropping systems in the US Corn Belt*, in: V. O. Sadras and D. F. Calderini (Eds.), *Crop physiology. Applications for genetic improvement and agronomy*, Oxford: Academic Press. pp. 15-42.
- Taylor H.M. (1980) *Soybean growth and yield as affected by row spacing and by seasonal water supply*. *Agron J* 72:543-547. DOI: 10.2134/agronj1980.00021962007200030032x.
- Wilhelm W.W., Wortmann C.S. (2004) *Tillage and rotation interactions for corn and soybean grain yield as affected by precipitation and air temperature*. *Agron J* 96:425-432. DOI: 10.2134/agronj2004.4250.

**Originally published:** Juan Ignacio Rattalino Edreira, S. Mourtzinis, S.P. Conley, A.C. Roth; I.A. Ciampitti, M. A. Licht, H. Kandel, P.M. Kyveryga, L.E. Lindsey, D.S. Mueller, S.L. Naeve, E. Nafziger, J.E. Specht, J. Stanley; M.J. Staton, P. Grassini. 2017. *Assessing causes of yield gaps in agricultural areas with diversity in climate and soils*. *Agricultural and Forest Meteorology*. <http://dx.doi.org/10.1016/j.agrformet.2017.07.010>